

QUESTION 6: What immune system-enhancing strategies can be employed to reduce the risk of surgical site infections/periprosthetic joint infections (SSIs/PJIs)?

RECOMMENDATION: Besides medical optimization of patients to enhance their immunity, there is some evidence demonstrating that immunonutrients (amino acids), vitamin D supplementation and passive/active immunization against *Staphylococcus aureus* may enhance immune system function, and potentially reduce the incidence of SSIs/PJIs.

LEVEL OF EVIDENCE: Moderate

DELEGATE VOTE: Agree: 74%, Disagree: 11%, Abstain: 15% (Super Majority, Strong Consensus)

RATIONALE

There is a close relationship between immunity and SSIs and PJIs. Thus, the strengthening of the immune system may reduce SSIs and PJIs. The strongest rationale for immune system enhancing strategies to reduce the risk of SSIs and PJIs is that perioperative immunosuppressive therapy is believed to increase these complications. This thinking has led to empirical bundles that include stopping immunosuppressive drugs (i.e., glucocorticoids, disease-modifying antirheumatic drugs (DMARDs) and biologic agents) before elective surgery [1]. Other investigators have concluded that while there is evidence to support the use of methotrexate perioperatively in rheumatoid arthritis patients, it remains unclear whether using anti-tumor necrosis factor (anti-TNF) medications perioperatively increases the risk of SSI [2].

Although cessation of immunosuppressive therapy prior to elective surgery has been adopted as a standard of care for the aforementioned reasons [3,4], there are no data from randomized, double-blind controlled clinical trials available to guide immunosuppressive therapy in the perioperative setting [5]. Thus, to identify the available information on this subject, a systematic review was completed on the peer-reviewed literature identified by a PubMed search performed on February 24, 2018 using the keywords “immunosuppression” or “immunostimulatory,” and “SSI” or “PJI” or “elective surgery.” This literature search identified 60 references from 1992 to 2018. After eliminating 49 that did not contain information directly addressing the question, the remaining 11 were divided into two categories: Primary Clinical Research (n = 7, four studies were positive [6–9] and three studies were negative [10–12]) and Clinical Reviews (n = 4, all reviews were positive [1,2,5,13]). Of note, a review of the pre-clinical literature failed to identify any research aimed at answering this question.

Activation of the immune system by active and passive immunization is a method that has been applied for many years to cope with many infective organisms. Recently, promising studies have been conducted on active and passive immunization for *Staphylococcus aureus*, which is the main causative agent identified for PJIs [14,15]. Although a vaccine for *S. aureus* has not been introduced clinically, a clinical trial by Pfizer is underway at the moment evaluating the effect of a tetravalent vaccine on patients undergoing spine surgery. There is also the potential for the development of a vaccine against *Pseudomonas* [16,17].

The relationship between immunity and nutrients has long been studied in patients with a poor immune system. The use of glutamine, arginine, omega-3 polyunsaturated fatty acids and ribonucleic acids in the perioperative period has been reported to reduce postoperative complications [18]. In a meta-analysis conducted by Zheng et al., 13 randomized controlled trials including 1,269 patients were evaluated. The meta-analysis revealed that the addition of immunonutrients to routine preoperative diets reduced subsequent SSIs and shortened the hospital stays [19]. Moreover, immunomodulator effects of Eicosapentaenoic acid (EPA) have been elucidated [19]. In a prospective study by Horie et al., administration of preoperative arginine-enriched nutrition reduced superficial, deep and organ-space infection in a cohort of patients undergoing colorectal cancer surgery [20]. On the other hand, one study found that preoperative or perioperative immunonutrition did not reduce the postoperative infectious complications and SSIs in head and neck cancer patients [10].

Vitamin D is an important immune system enhancer, playing an essential role in neutrophil motility, activation of macrophages and inducing T-helper type 1 cells, which target bacterial pathogens that are commonly responsible for PJIs [21,22]. A recent study by Traven et al. demonstrated that low-serum vitamin D levels (25-OH) in patients undergoing joint arthroplasty were associated with an increased risk of 90-day complications as well as PJIs [23]. However, to date, no studies exist to demonstrate that correction of vitamin D deficiency repudiates the reported association. In addition, it is not known what dose and duration of vitamin D supplement are required to correct the deficiency.

Vitamin E also plays an important role in enhancing immune system function via its antioxidant properties. It also reduces apoptosis and increases macrophage activation. Chen et al. demonstrated that murine macrophages with vitamin E-enriched ultra-high molecular weight polyethylene (VE-UHMWPE) particles induced less apoptosis and Tumor Necrosis Factor (TNF) release versus particles without vitamin E [24]. Banche et al. demonstrated that VE-UHMWPE provides a less adhesive surface to *S. aureus* and *E. coli* [25]. On the other hand, Williams et al. reported that the addition of vitamin E to UHMWPE might not reduce clinically relevant rates of biofilm-related PJIs [26]. Further studies are required to better delineate the role of vitamin E in preventing PJIs.

The relationship between smoking and immunity has been established [27]. Smoking, in particular, causes immunosuppression by inactivating macrophages, neutrophils, natural killer cells and lymphocytes [27]. Moreover, smoking causes tissue hypoxia and slows blood flow to tissues potentially preventing the immune cells to reach infecting organisms in a given tissue. Smoking cessation is likely to restore immune function and potentially minimize the risk of subsequent SSIs/PJIs [28].

Greenky et al. have shown that patients with preoperative anemia (hemoglobin level less than 13 g/dL in men and 12 g/dL in women) are at greater risk of PJIs (4.3% in anemic patients compared with 2% in non-anemic patients) [29]. The association between anemia and a higher rate of SSI/PJI may be explained by numerous factors. Patients with anemia are more likely to have tissue hypoxia, which adversely affects wound healing. Patients with anemia may suffer chronic conditions such as renal disease that in their own right may be associated with SSIs/PJIs. Patients with anemia may be subjected to a higher rate of allogeneic blood transfusion with its immunomodulating effects.

Another cause of immunosuppression is malnutrition. Bohl et al. reported that patients with hypoalbuminemia are at a greater risk of developing PJIs following joint arthroplasty [30]. Malnutrition can be defined as a serum albumin level < 3.5 g/dL, serum transferrin levels < 200 mg/dL, serum prealbumin < 15 gm/dL, and total lymphocyte count (TLC) < 1,500 cells/mm³ [31]. Dialysis therapy due to renal insufficiency, chronic hepatic insufficiency, malnutrition and depression-psychosis may cause hypoalbuminemia [32]. We should state that the current definitions of malnutrition mostly concentrate on protein deficiency, and the importance of other nutritional parameters such as vitamins, minerals, etc. are not well-studied.

This literature review also found evidence of nonspecific global health treatments that have been described as being immune system enhancing to reduce SSIs/PJIs. These include maintaining body temperature, high concentration of oxygen [13], perioperative glucose control [9] and eliminating blood transfusions [6].

With the available evidence, it is reasonable to propose that discontinuation of immunosuppressive agents, medical optimization of patients with chronic conditions, such as anemia and diabetes, and administration of immunonutrients, such as amino acids and vitamins, are likely to lead to better outcomes after surgical procedures in general and a reduced rate of SSIs and PJIs in particular. Future studies will reveal if vaccines against organisms such as *Staphylococcus aureus* are effective in reducing the incidence of SSIs/PJIs after orthopaedic and other surgical procedures.

REFERENCES

- [1] Härle P, Straub RH, Fleck M. Elective surgery in rheumatic disease and immunosuppression: to pause or not. *Rheumatology (Oxford)*. 2010;49:1799–1800. doi:10.1093/rheumatology/keq049.
- [2] Morrison TA, Figgie M, Miller AO, Goodman SM. Periprosthetic joint infection in patients with inflammatory joint disease: a review of risk factors and current approaches to diagnosis and management. *HSS J*. 2013;9:183–194. doi:10.1007/s11420-013-9338-8.
- [3] Rogers SO. Surgical perspective: centers for disease control and prevention guideline for the prevention of surgical site infection 2017. *Surg Infect (Larchmt)*. 2017;18:383–384. doi:10.1089/sur.2017.097.
- [4] Berríos-Torres SI, Umscheid CA, Bratzler DW, Leas B, Stone EC, Kelz RR, et al. Centers for disease control and prevention guideline for the prevention of surgical site infection, 2017. *JAMA Surg*. 2017;152:784–791. doi:10.1001/jamasurg.2017.0904.
- [5] Härle P, Straub RH, Fleck M. Perioperative management of immunosuppression in rheumatic diseases—what to do? *Rheumatol Int*. 2010;30:999–1004. doi:10.1007/s00296-009-1323-7.
- [6] Fragkou PC, Torrance HD, Pearse RM, Ackland GL, Prowle JR, Owen HC, et al. Perioperative blood transfusion is associated with a gene transcription profile characteristic of immunosuppression: a prospective cohort study. *Crit Care*. 2014;18:541. doi:10.1186/s13054-014-0541-x.
- [7] Ott E, Bange FC, Sohr D, Teebken O, Mattner F. Risk factors associated with surgical site infections following vascular surgery at a German university hospital. *Epidemiol Infect*. 2013;141:1207–1213. doi:10.1017/S095026881200180X.
- [8] Berbari EF, Osmon DR, Lahr B, Eckel-Passow JE, Saras G, Hanssen AD, et al. The Mayo prosthetic joint infection risk score: implication for surgical site infection reporting and risk stratification. *Infect Control Hosp Epidemiol*. 2012;33:774–781. doi:10.1086/666641.
- [9] Sehgal R, Berg A, Figueroa R, Poritz LS, McKenna KJ, Stewart DB, et al. Risk factors for surgical site infections after colorectal resection in diabetic patients. *J Am Coll Surg*. 2011;212:29–34. doi:10.1016/j.jamcollsurg.2010.09.011.
- [10] Falewee MN, Schilf A, Boufflers E, Cartier C, Bachmann P, Pressoir M, et al. Reduced infections with perioperative immunonutrition in head and neck cancer: exploratory results of a multicenter, prospective, randomized, double-blind study. *Clin Nutr*. 2014;33:776–784. doi:10.1016/j.clnu.2013.10.006.
- [11] Everhart JS, Altneu E, Calhoun JH. Medical comorbidities are independent preoperative risk factors for surgical infection after total joint arthroplasty. *Clin Orthop Relat Res*. 2013;471:3112–3119. doi:10.1007/s11999-013-2923-9.
- [12] Dahl RM, Wetterslev J, Jorgensen LN, Rasmussen LS, Moller AM, Meyhoff CS, et al. The association of perioperative dexamethasone, smoking and alcohol abuse with wound complications after laparotomy. *Acta Anaesthesiol Scand*. 2014;58:352–361. doi:10.1111/aas.12270.
- [13] Kawasaki T, Sata T. Perioperative innate immunity and its modulation. *J UOEH*. 2011;33:123–137.
- [14] Sjøe NH, Jensen NV, Jensen AL, Koch J, Poulsen SS, Pier GB, et al. Active and passive immunization against staphylococcus aureus periprosthetic osteomyelitis in rats. *In Vivo*. 2017;31:45–50. doi:10.21873/invivo.11023.
- [15] Gustin M-P, Ohannessian R, Giard M, Caillat-Vallet E, Savay A, Vanhems P, et al. Use of surveillance data to calculate the sample size and the statistical power of randomized clinical trials testing staphylococcus aureus vaccine efficacy in orthopedic surgery. *Vaccine*. 2017;35:6934–6937. doi:10.1016/j.vaccine.2017.10.068.
- [16] de Bruyn G, Saleh J, Workman D, Pollak R, Elinoff V, Fraser NJ, et al. Defining the optimal formulation and schedule of a candidate toxoid vaccine against *Clostridium difficile* infection: A randomized Phase 2 clinical trial. *Vaccine*. 2016;34:2170–2178. doi:10.1016/j.vaccine.2016.03.028.
- [17] Döring G, Meisner C, Stern M, Flagella Vaccine Trial Study Group. A double-blind randomized placebo-controlled phase III study of a *Pseudomonas aeruginosa* flagella vaccine in cystic fibrosis patients. *Proc Natl Acad Sci*. 2007;104:11020–11025. doi:10.1073/pnas.0702403104.
- [18] Ryan AM, Reynolds JV, Healy L, Byrne M, Moore J, Brannelly N, et al. Enteral nutrition enriched with eicosapentaenoic acid (EPA) preserves lean body mass following esophageal cancer surgery: results of a double-blinded randomized controlled trial. *Ann Surg*. 2009;249:355–363. doi:10.1097/SLA.0b013e31819a4789.

- [19] Zheng Y, Li F, Qi B, Luo B, Sun H, Liu S, et al. Application of perioperative immunonutrition for gastrointestinal surgery: a meta-analysis of randomized controlled trials. *Asia Pac J Clin Nutr*. 2007;16 Suppl 1:253–257.
- [20] Horie H, Okada M, Kojima M, Nagai H. Favorable effects of preoperative enteral immunonutrition on a surgical site infection in patients with colorectal cancer without malnutrition. *Surg Today*. 2006;36:1063–1068. doi:10.1007/s00595-006-3320-8.
- [21] Rode AKO, Kongsbak M, Hansen MM, Lopez DV, Levring TB, Woetmann A, et al. Vitamin D counteracts mycobacterium tuberculosis-induced cathelicidin downregulation in dendritic cells and allows Th1 differentiation and IFN γ secretion. *Front Immunol*. 2017;8:656. doi:10.3389/fimmu.2017.00656.
- [22] Hewison M. Vitamin D and the immune system: new perspectives on an old theme. *Endocrinol Metab Clin North Am*. 2010;39:365–379, table of contents. doi:10.1016/j.ecl.2010.02.010.
- [23] Traven SA, Chiaramonti AM, Barfield WR, Kirkland PA, Demos HA, Schutte HD, et al. Fewer complications following revision hip and knee arthroplasty in patients with normal vitamin D levels. *J Arthroplasty*. 2017;32:S193–S196. doi:10.1016/j.arth.2017.02.038.
- [24] Chen W, Bichara DA, Suhardi J, Sheng P, Muratoglu OK. Effects of vitamin E-diffused highly cross-linked UHMWPE particles on inflammation, apoptosis and immune response against *S. aureus*. *Biomaterials*. 2017;143:46–56. doi:10.1016/j.biomaterials.2017.07.028.
- [25] Banche G, Allizond V, Bracco P, Bistolfi A, Boffano M, Cimino A, et al. Interplay between surface properties of standard, vitamin E blended and oxidised ultra high molecular weight polyethylene used in total joint replacement and adhesion of staphylococcus aureus and escherichia coli. *Bone Joint J*. 2014;96-B:497–501. doi:10.1302/0301-620X.96B4/32895.
- [26] Williams DL, Vinciguerra J, Lerdahl JM, Bloebaum RD. Does vitamin E-blended UHMWPE prevent biofilm formation? *Clin Orthop Relat Res*. 2015;473:928–935. doi:10.1007/s11999-014-3673-z.
- [27] Springer BD. Modifying risk factors for total joint arthroplasty: strategies that work nicotine. *J Arthroplasty*. 2016;31:1628–1630. doi:10.1016/j.arth.2016.01.071.
- [28] Bedard NA, Dowdle SB, Owens JM, Duchman KR, Gao Y, Callaghan JJ. What is the impact of smoking on revision total hip arthroplasty? *J Arthroplasty*. 2018;33:S182–S185. doi:10.1016/j.arth.2017.12.041.
- [29] Greenky M, Gandhi K, Pulido L, Restrepo C, Parvizi J. Preoperative anemia in total joint arthroplasty: is it associated with periprosthetic joint infection? *Clin Orthop Relat Res*. 2012;470:2695–2701. doi:10.1007/s11999-012-2435-z.
- [30] Bohl DD, Shen MR, Kayupov E, Cvetanovich GL, Della Valle CJ. Is hypoalbuminemia associated with septic failure and acute infection after revision total joint arthroplasty? A study of 4517 patients from the national surgical quality improvement program. *J Arthroplasty*. 2016;31:963–967. doi:10.1016/j.arth.2015.11.025.
- [31] Morey VM, Song YD, Whang JS, Kang YG, Kim TK. Can serum albumin level and total lymphocyte count be surrogates for malnutrition to predict wound complications after total knee arthroplasty? *J Arthroplasty*. 2016;31:1317–1321. doi:10.1016/j.arth.2015.12.004.
- [32] Aldebeyan S, Nooh A, Aoude A, Weber MH, Harvey EJ. Hypoalbuminaemia—a marker of malnutrition and predictor of postoperative complications and mortality after hip fractures. *Injury*. 2017;48:436–440. doi:10.1016/j.injury.2016.12.016.